

Influence of Gaps in Earth Radiation Budget Climate Data Records

Norman Loeb, Bruce Wielicki, Tak Wong and Peter Parker
NASA LaRC


CERES Science Team Meeting
NASA GISS, New York, Oct 27-29, 2008

Introduction


- How does a gap in a climate data record affect our ability to constrain cloud feedback?
- How much overlap is needed between successive CERES instruments?

Tropical Mean (20°N to 20°S) Outgoing Longwave Radiation

Without Overlap Adjustment




With Overlap Adjustment



- *Instrument-to-instrument absolute calibration differences are 1 to 4 Wm^{-2} .*
- *=> Absolute accuracy alone is insufficient to detect climate change at the 0.6 Wm^{-2} per decade level of anthropogenic radiative forcing by greenhouse gases.*
- *Overlapping observations allows the use of instrument stability instead of absolute accuracy to constrain decadal climate change.*

Radiation Budget Gap Risk: Satellite Scenarios

Past and Current Scenarios for NPP, NPOESS



CERES Climate Data Record Observational Requirements for Constraining Cloud Feedback

- The largest uncertainty in global climate sensitivity over the next century is cloud feedback (especially for low clouds).
- A global cloud feedback of 25% would amplify or dampen global warming by 25%.
- CERES can observe decadal changes in cloud radiative forcing that constrain the large uncertainty in cloud feedback and therefore climate sensitivity.
- Uncertainty in the CRE^{net} trend should be less **than 0.15 Wm^{-2} per decade** in order to constrain cloud radiative feedback to 25% of the anticipated change in anthropogenic radiative forcing over the next few decades (0.6 Wm^{-2} ; Houghton et al. 2001).

Simulated 30-Yr Record in Net Cloud Radiative Effect (Based on First 5 Yr of CERES-Terra Obs)




Figure 1 One realization of a 30-year time series in net cloud radiative effect for (a) no gap; (b) one-year gap starting in 2014 with an absolute calibration shift of -2% in the SW and -1% in the LW across the gap. Solid lines are least-square linear fits.

Uncertainty in net Cloud Radiative Effect due to Absolute Calibration Uncertainty

$$CRE^{sw} = F_{clr}^{sw} - F_{all}^{sw}$$

$$CRE^{lw} = F_{clr}^{lw} - F_{all}^{lw}$$

$$CRE^{net} = CRE^{sw} + CRE^{lw}$$

$$\delta CRE^X = \sqrt{(\delta F_{clr}^X)^2 + (\delta F_{all}^X)^2 - 2\rho^X \delta F_{clr}^X \delta F_{all}^X}$$

$$\delta F_{clr}^X = \Delta G F_{clr}^X, \quad \delta F_{all}^X = \Delta G F_{all}^X$$

$$\delta CRE^{net} = \sqrt{(\delta CRE^{sw})^2 + (\delta CRE^{lw})^2 + 2\rho^{net} \delta CRE^{sw} \delta CRE^{lw}}$$

Trend Error Due to a Gap in Record

$$E(\hat{\omega}) = \sqrt{(\hat{\omega} - \omega)^2 + (\delta\hat{\omega} - \delta\omega)^2}$$

ω Trend (no gap)


$\hat{\omega}$ Trend (with gap)

$\delta\omega$ 95% Confidence interval in trend (no gap)


$\delta\hat{\omega}$ 95% Confidence interval in trend (with gap)

Error (2σ) in Net Cloud Radiative Effect Trend Due to 1-Yr Gap as a Function of SW and LW Absolute Calibration


Gap at Beginning of Record



Gap in Middle of Record



How Much Overlap Time is Necessary Between Successive Instruments?




$$Z = | \langle X \rangle - \langle Y \rangle |$$

$$\delta Z_j = \sqrt{\delta X_j^2 + \delta Y_j^2 - 2\rho_{X_j Y_j} \delta X_j \delta Y_j}$$

Determine L such that $\delta Z < 0.15 \text{ Wm}^{-2} \text{ decade}^{-1}$

- Use one year of overlap between CERES Terra & Aqua to determine L.


Uncertainty in Terra–Aqua Monthly Mean Difference in CRE



- Determine monthly mean CRE from daily means for CERES Terra and Aqua.
- For each month compute $\delta\text{CRE}(\text{Terra})$, $\delta\text{CRE}(\text{Aqua})$ and ρ .
- Above plot shows uncertainty in the Terra-Aqua CRE difference for each month and when several months are combined.

Overlap Requirements

- Determine number of overlapping months required to reduce random uncertainty in the difference between the two overlapping data records to ensure a 30-year trend uncertainty of $< 0.15 \text{ Wm}^{-2}$ per decade.
- Use overlap between Terra and Aqua in 2004 to determine random error.



Multiple overlapping months: the trend uncertainty reaches 0.15 Wm^{-2} per decade after ~ 3 months for the globe, and 6 months for the tropics.

Conclusions

- Current ERB instruments are far more precise than they are absolutely accurate.
- A gap in the middle of the data record has far greater impact than a gap near the beginning.
- To constrain cloud radiative feedback to 25% of anthropogenic forcing in next few decades, the absolute calibration change across the gap must be $< 0.3\%$ in the SW and 0.1% in the LW.
 - => This is well beyond the capability of current ERB instruments. A gap of any length restarts the record at zero.
- To overcome the effect of a gap, observing systems require six months of overlap between successive instruments in order to tie the data record of one instrument to the next.
 - => More details in Loeb et al., JGR, 2008 (submitted)

CERES Flight Schedule

Enabling Climate Data Record Continuity

Spacecraft	Instruments	Launch	Science Initiation	Collected Data (Months)
TRMM	PFM	11/97	1/98	9
Terra	FM1, FM2	12/99	3/00	98 +
Aqua	FM3, FM4	5/02	6/02	71 +
NPP	FM5	June 2010	-	-
<i>NPOESS C1</i>	<i>FM6</i>	<i>January 2013</i>	-	-
<i>NPOESS C3</i>	<i>CERES follow-on</i>	<i>January 2018</i>	-	-

31 Instrument Years of Data